

Down Syndrome Healthcare Guidelines (2011 Revision) Record Sheet*

	Birth	6 mo	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
Genetic Counseling ¹ , Karyotype																							
Cardiology	Echo ⁴																						
CBC to R/O transient myeloproliferative disorder, polycythemia																							
Swallowing assessment if feeding problems or aspiration																							
Parent Group Info and Support	Parent-to-parent contact, support groups, current books and pamphlets																						
Hemoglobin			Hemoglobin annually beginning at 1 year old. If Hg<11, do (a) CRP and ferritin, or (b) Reticulocyte Hemoglobin Content (CHr). If possible risk for iron deficiency, do (a) or (b) regardless.																				
23-valent pneumococcal vaccine ²																							
Audiological Evaluation	ABR or OAE	Every 6 months until 3 years of age																					
Ophthalmologic Evaluation	Red Reflex	Ophth Appt	Annual ophthalmology appt					Q2 Ophthalmology appointment					Q3 Ophthalmology appointment										
Celiac Disease Screening					(Only test if signs and symptoms present)																		
Thyroid – TSH, T4	State Screen	Lab	Labs 12mo & 18mo	Test TSH and T4 annually																			
Neck X-ray (AAI) ³				✓ ⁴																			
Dental Exam			Annual Dental Exams. Reassure parents that delayed or irregular eruption, hypodontia are common.																				
Sleep Study by age 4 years	Done prior to 4 years of age																						
Early Intervention																							
Childhood					Discuss self-help, ADHD, OCD, wandering off, transition to middle school																		
Puberty																							
Facilitate transition																							
Sexual development and behaviors																							
Preventive care	Annually monitor for signs and symptoms of constipation, OSA, and aspiration.																						

1. Discuss Recurrence Rate of future pregnancies with parents
2. 23-valent pneumococcal vaccine if chronic or pulmonary disease.
3. AAI: See AAP Guidelines page 399 - X-rays only if myopathic signs or symptoms * *Peds* 2011 ;128 :393-406 Chart by Sie Center for Down Syndrome
4. Follow up to be determined by Cardiologist